

Using Aura MLS observations to model budget terms, strattrop fluxes, and surface variability of N₂O

Daniel Ruiz & Michael Prather,

Earth System Science Dept, UC Irvine

N₂O loss from Aura MLS measurements







Identifier	Version	Spectral Region	Availability	Comments
N2O-640-V3	V3.3	640 GHz	08/2004-12/2010	Prather et al. 2015
N2O-640-V4	V4.2x	640 GHz	08/2004-12/2013	MLS instrument issue; channel data not recommended after 07/2013
N2O-190-V4	V4.2x	190 GHz	08/2004-12/2018	Retrieval bias for profiles >68 hPa, not used

$N_2O - MIP$ (UCI CTM, NASA GMI, NILU LMDZ5)


Decay per N-tracers are simulated in the absence of surface emissions to preserve the stratospheric signal



Measurements vs Models




MLS N₂O* uses NOAA HATS N₂O burden for 2005-2018 (ppb)



MLS N_2O^* uses NOAA HATS N_2O burden for 2005-2018 (ppb) UCI F11x* decay corresponds to the right axis. All others correspond to the left.

Complementary N2O (Neg-N2O) flux vs N2O(x) loss


Stratospheric N₂O loss leads to peak neg-N₂O flux ~1 year later



X-correlation w/ cN2Ox STE	Offset
MLS N2O loss	-1.0 yrs
UCI N2Ox loss (MLS overlap period)	-0.9 yrs
UCI N2Ox loss (entire period)	-1.0 yrs


cN₂Ox: every N₂Ox molecule lost produces one cN2Ox molecule

Monthly Tropospheric N2Ox vs total cN2Ox STE flux




X-correlation w/ cN2Ox STE	Offset
UCI N2Ox Northern Hemisphere	+0.2 yrs
UCI N2Ox Southern Hemisphere	+0.7 yrs

Surface N₂Ox: Seasonal cycle of stratospheric loss




- North Larger seasonal amplitude
- South < North

Monthly surface climatologies



UCI N₂Ox (no emissions) agrees with NOAA NH seasonal cycle

UCI CTM STE: How do the STE fluxes compare in each




negative-N₂O STE SH > NH (mechanism not understood)

N₂O vs O₃ Tracer Correlations:

Lower N₂Ox (no emissions) in SH for same O₃ 2.5 **UCI CTM** SH = -0.018x + 4.82.0 O_3 (ppm) 0.5 0.0 160 180 200 220 240 260 N₂Ox (ppb)

 Hemispheric difference due to increased transport of depleted-N₂O air into SH?



*Data > 70° were not used due to retrieval limitations of ACE orbit

Acknowledgements





- Jet Propulsion Laboratory Lucien Froidevaux
- GSFC Susan Strahan, Steve Steenrod
- NILU Rona Thompson
- NOAA James Elkins, Geoff Dutton
- NASA Ken Jucks